A1
Hart District Local Plan
Queen's Road, North Warnborough
Agricultural Land Classification
ALC Map and Report
May 1995

# AGRICULTURAL LAND CLASSIFICATION REPORT

# HART DISTRICT LOCAL PLAN QUEEN'S ROAD, NORTH WARNBOROUGH

## 1. Summary

- 1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality for a number of sites in the Hart district of Hampshire. This forms part of MAFF's input to the preparation of the Hart District Local Plan.
- 1.2 Approximately 5 hectares of land south of Queen's Road and to the east of Robert May's School in North Warnborough, Hampshire, was surveyed during May 1995. The Agricultural Land Classification (ALC) survey was undertaken at a detailed level of approximately one boring per hectare. A total of 4 auger borings and 2 soil inspection pits were assessed in accordance with MAFF's revised guidelines and criteria for grading the quality of agricultural land (MAFF, 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose long-term limitations on its use for agriculture.
- 1.3 The work was carried out by members of the Resource Planning Team in the Guildford Statutory Group of ADAS.
- 1.4 At the time of the survey all of the agricultural land on this site was under arable cropping. The Non-Agricultural land mapped comprises a public footpath.
- 1.5 The distribution of grades and subgrades is shown on the attached ALC map and the areas are given in Table 1 below. The map has been drawn at a scale of 1:10,000. It is accurate at this scale, but any enlargement would be misleading. This map supersedes any previous ALC information for this site.

Table 1: Distribution of Grades and Subgrades

| Grade              | Area (ha)  | % of Site  |
|--------------------|------------|------------|
| 2                  | 4.4        | 93.6       |
| Non-Agricultural   | <u>0.3</u> | <u>6.4</u> |
| Total area of site | 4.7        | 100%       |

- 1.6 Appendix I gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.
- 1.7 All of the agricultural land on this site has been classified as Grade 2, very good quality, the key limitation being soil droughtiness. The soils are derived from the Upper Chalk and as such comprise slightly stony, calcareous, silty clay loams and clays over Chalk at depth These soils experience a limitation on the amount of water that is available for use by crops.

#### 2. Climate

- 2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.
- 2.2 The main parameters used in the assessment of an overall climatic limitation are average annual rainfall, as a measure of overall wetness, and accumulated temperature (day degrees Celsius, Jan-June), as a measure of the relative warmth of a locality.
- 2.3 A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met. Office, 1989). The details are given in the table below and these show that there is no overall climatic limitation affecting the site. However, climatic factors do interact with soil properties to influence soil wetness and droughtiness limitations.
- 2.4 No local climatic factors such as exposure or frost risk are believed to affect the site.

**Table 2: Climatic Interpolations** 

| Grid Reference                  | SU 734 512 |
|---------------------------------|------------|
| Altitude (m)                    | 85         |
| Accumulated Temperature         | 1435       |
| (day degrees, Jan-June)         |            |
| Average Annual Rainfall (mm)    | 739        |
| Field Capacity (days)           | 158        |
| Moisture Deficit, Wheat (mm)    | 105        |
| Moisture Deficit, Potatoes (mm) | 98         |
| Overall Climatic Grade          | 1          |

#### 3. Relief

3.1 The land on this site slopes gently from approximately 90m AOD in the south east to 85m AOD in the north west. Nowhere on the site does altitude or relief impose limitations to agricultural land quality.

### 4. Geology and Soil

- 4.1 The relevant geological sheet (BGS, 1981) maps the entire site as Upper Chalk.
- 4.2 The published soil information for the site (SSEW, 1983) shows the Andover 1 association across the site. These soils are described as 'Shallow well drained calcareous silty soils over chalk on slopes and crests. Deep calcareous and non-calcareous fine silty soils in valley bottoms. Striped soil pattern locally.' (SSEW, 1983).
- 4.3 Detailed field survey broadly confirms the existence of soils similar to those described in paragraph 4.2 with slightly heavier soils on the hill top.

#### 5. Agricultural Land Classification

- 5.1 Table 1 provides the details of the area measurements for each grade and the distribution of each grade is shown on the attached ALC map.
- 5.2 The location of the soil observation points are shown on the attached sample point map.

#### Grade 2

- Pit 1 is representative of the majority of profiles which comprise very slightly to slightly 5.3 flinty (3-8% total stone & 1-2% > 2cm), calcareous, medium silty clay loam topsoils over moderately structured upper subsoils of similar texture. The flint content in the upper subsoil measures 5%, however, above the chalk bedrock this increases to 25% flint in some profiles and 50% chalk in others. The Upper Chalk occurs at 55-88cm from the surface but plant roots are able to penetrate, and thus extract water from, a further 20cm into the chalk. The amount of flint and chalk fragments present within the profile, together with the slightly retricted depth of soil over chalk, combine to reduce the amount of profile available water for crops. This land is therefore subject to a minor soil droughtiness limitation.
- 5.4 In the south east of the site, on the hill top, the profiles are slightly heavier in texture. The topsoil is similar to those across the rest of the site but the upper subsoils comprise very slightly flinty (2% total flint) heavy silty clay loams and clays. Pit 2 showed the heavy subsoils to be moderately structured and therefore well drained. At approximately 65cm depth the Upper Chalk is encountered which shows evidence of plant roots to 85cm depth. This land is therefore also limited to Grade 2 by soil droughtiness.

ADAS Ref: 1506/41/95 MAFF Ref: 15/347

Resource Planning Team Guildford Statutory Group

ADAS Reading

# **SOURCES OF REFERENCE**

British Geological Survey (1981): Sheet No. 284, Basingstoke, 1:50,000 Scale (solid & drift edition).

MAFF (1988), Agricultural Land Classification of England and Wales: Revised guidelines and criteria for grading the quality of agricultural land.

Meteorological Office (1989), Climatological Data for Agricultural Land Classification.

Soil Survey of England and Wales (1983), Sheet 6, Soils of South East England, and accompanying legend.

#### APPENDIX I

#### DESCRIPTION OF THE GRADES AND SUBGRADES

### Grade 1: Excellent Quality Agricultural Land

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

# Grade 2: Very Good Quality Agricultural Land

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land of this grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1 land.

# Grade 3: Good to Moderate Quality Land

Land with moderate limitations which affect the choice of crops, the timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown, yields are generally lower or more variable than on land in Grades 1 and 2.

## Subgrade 3a: Good Quality Agricultural Land

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

# Subgrade 3b: Moderate Quality Agricultural Land

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

### Grade 4: Poor Quality Agricultural Land

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (eg. cereals and forage crops) the yields of which are variable. In moist climates, yields of grass may be moderate to high but there may be difficulties in utilisation. The grade also includes very droughty arable land.

#### Grade 5: Very Poor Quality Agricultural Land

Land with severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

#### Urban

Built-up or 'hard' uses with relatively little potential for a return to agriculture including: housing, industry, commerce, education, transport, religous buildings, cemetries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

# Non-agricultural

'Soft' uses where most of the land could be returned relatively easily to agriculture, including: private parkland, public open spaces, sports fields, allotments and soft-surfaced areas on airports. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

#### Woodland

Includes commercial and non-commercial woodland. A distinction may be made as necessary between farm and non-farm woodland.

# Agricultural Buildings

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (eg. polythene tunnels erected for lambing) may be ignored.

# Open Water

Includes lakes, ponds and rivers as map scale permits.

### Land Not Surveyed

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above, eg. buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will be shown.

# APPENDIX II

# FIELD ASSESSMENT OF SOIL WETNESS CLASS

### SOIL WETNESS CLASSIFICATION

Soil wetness is classified according to the depth and duration of waterlogging in the soil profile. Six soil wetness classes are identified and are defined in the table below.

### **Definition of Soil Wetness Classes**

| Wetness Class | Duration of Waterlogging <sup>1</sup>   |
|---------------|---|
| I             | The soil profile is not wet within 70 cm depth for more than 30 days in most years. <sup>2</sup>  |
| п             | The soil profile is wet within 70 cm depth for 31-90 days in most years or, if there is no slowly permeable layer within 80 cm depth, it is wet within 70 cm for more than 90 days, but only wet within 40 cm depth for 30 days in most years.                          |
| m             | The soil profile is wet within 70 cm depth for 91-180 days in most years or, if there is no slowly permeable layer present within 80 cm depth, it is wet within 70 cm for more than 180 days, but only wet within 40 cm depth for between 31-90 days in most years.     |
| IV            | The soil profile is wet within 70 cm depth for more than 180 days but not wet within 40 cm depth for more than 210 days in most years or, if there is no slowly permeable layer present within 80 cm depth, it is wet within 40 cm depth for 91-210 days in most years. |
| v             | The soil profile is wet within 40 cm depth for 211-335 days in most years.  |
| VI            | The soil profile is wet within 40 cm depth for more than 335 days in most years.  |

Soils can be allocated to a wetness class on the basis of quantitative data recorded over a period of many years or by the interpretation of soil profile characteristics, site and climatic factors. Adequate quantitative data will rarely be available for ALC surveys and therefore the interpretative method of field assessment is used to identify soil wetness class in the field. The method adopted here is common to ADAS and the SSLRC.

<sup>&</sup>lt;sup>1</sup>The number of days specified is not necessarily a continuous period.

<sup>&</sup>lt;sup>2</sup>'In most years' is defined as more than 10 out of 20 years.

# **APPENDIX III**

# SOIL PIT AND SOIL BORING DESCRIPTIONS

# Contents:

Soil Abbreviations - Explanatory Note

Soil Pit Descriptions

**Database Printout - Boring Level Information** 

**Database Printout - Horizon Level Information** 

### SOIL PROFILE DESCRIPTIONS: EXPLANATORY NOTE

Soil pit and auger boring information collected during ALC fieldwork is held on a computer database. This uses notations and abbreviations as set out below.

# **Boring Header Information**

- 1. **GRID REF**: national 100 km grid square and 8 figure grid reference.
- 2. USE: Land use at the time of survey. The following abbreviations are used.

ARA: Arable WHT: Wheat BAR: Barley
CER: Cereals OAT: Oats MZE: Maize
OSR: Oilseed rape BEN: Field Beans BRA: Brassicae
POT: Potatoes SBT: Sugar Beet FCD: Fodder Crops

LIN: Linseed FRT: Soft and Top Fruit FLW: Fallow

PGR: Permanent PastureLEY: Ley Grass RGR: Rough Grazing SCR: Scrub CFW: Coniferous Woodland DCW: Deciduous Wood

HTH: Heathland BOG: Bog or Marsh FLW: Fallow PLO: Ploughed SAS: Set aside OTH: Other

HRT: Horticultural Crops

- 3. GRDNT: Gradient as estimated or measured by a hand-held optical clinometer.
- 4. GLEY/SPL: Depth in centimetres (cm) to gleying and/or slowly permeable layers.
- 5. AP (WHEAT/POTS): Crop-adjusted available water capacity.
- 6. MB (WHEAT/POTS): Moisture Balance. (Crop adjusted AP crop adjusted MD)
- 7. **DRT**: Best grade according to soil droughtiness.
- 8. If any of the following factors are considered significant, 'Y' will be entered in the relevant column.

MREL: Microrelief limitation FLOOD: Flood risk EROSN: Soil erosion risk EXP: Exposure limitation FROST: Frost prone DIST: Disturbed land

**CHEM**: Chemical limitation

9. LIMIT: The main limitation to land quality. The following abbreviations are used.

OC: Overall Climate AE: Aspect EX: Exposure FR: Frost Risk GR: Gradient MR: Microrelief FL: Flood Risk TX: Topsoil Texture DP: Soil Depth CH: Chemical WE: Wetness WK: Workability

**DR**: Drought **ER**: Erosion Risk **WD**: Soil Wetness/Droughtiness

ST: Topsoil Stoniness

# Soil Pits and Auger Borings

1. **TEXTURE**: soil texture classes are denoted by the following abbreviations.

S: Sand LS: Loamy Sand SL: Sandy Loam SZL: Sandy Silt Loam CL: Clay Loam ZCL: Silty Clay Loam

ZL: Silt Loam SCL: Sandy Clay Loam C: Clay

SC: Sandy Clay ZC: Silty Clay OL: Organic Loam **P**: Peat SP: Sandy Peat LP: Loamy Peat MZ: Marine Light Silts PL: Peaty Loam PS: Peaty Sand

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction will be indicated by the use of the following prefixes:

F: Fine (more than 66% of the sand less than 0.2mm)

M: Medium (less than 66% fine sand and less than 33% coarse sand)

C: Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content: M: Medium (<27% clay) H: Heavy (27-35% clay)

- 2. MOTTLE COL: Mottle colour using Munsell notation.
- 3. MOTTLE ABUN: Mottle abundance, expressed as a percentage of the matrix or surface described.

F: few <2% C: common 2-20% M: many 20-40% VM: very many 40% +

4. **MOTTLE CONT**: Mottle contrast

F: faint - indistinct mottles, evident only on close inspection

**D**: distinct - mottles are readily seen

P: prominent - mottling is conspicuous and one of the outstanding features of the horizon

- 5. **PED. COL**: Ped face colour using Munsell notation.
- 6. GLEY: If the soil horizon is gleyed a 'Y' will appear in this column. If slightly gleyed, an 'S' will appear.
- 7. STONE LITH: Stone Lithology One of the following is used.

HR: all hard rocks and stones SLST: soft oolitic or dolimitic limestone

CH: chalk FSST: soft, fine grained sandstone

**ZR**: soft, argillaceous, or silty rocks **GH**: gravel with non-porous (hard) stones

MSST : soft, medium grained sandstone GS : gravel with porous (soft) stones

This feet to a street a six best diese

SI: soft weathered igneous/metamorphic rock

Stone contents (>2cm, >6cm and total) are given in percentages (by volume).

8. STRUCT: the degree of development, size and shape of soil peds are described using the following notation:

<u>degree of development</u> WK: weakly developed MD: moderately developed

ST: strongly developed

ped size F: fine M: medium

C: coarse VC: very coarse

ped shape S: single grain M: massive

GR: granular AB: angular blocky

SAB: sub-angular blocky PR: prismatic

**PL**: platy

9. **CONSIST**: Soil consistence is described using the following notation:

L: loose VF: very friable FR: friable FM: firm VM: very firm

EM: extremely firm EH: extremely hard

10. SUBS STR: Subsoil structural condition recorded for the purpose of calculating profile droughtiness: G: good M: moderate P: poor

- 11. POR: Soil porosity. If a soil horizon has less than 0.5% biopores >0.5 mm, a 'Y' will appear in this column.
- 12. IMP: If the profile is impenetrable to rooting a 'Y' will appear in this column at the appropriate horizon.
- 13. SPL: Slowly permeable layer. If the soil horizon is slowly permeable a 'Y' will appear in this column.
- 14. CALC: If the soil horizon is calcareous, a 'Y' will appear in this column.
- 15. Other notations

APW: available water capacity (in mm) adjusted for wheat

APP: available water capacity (in mm) adjusted for potatoes

MBW: moisture balance, wheat MBP: moisture balance, potatoes

#### SOIL PIT DESCRIPTION

Site Name: HART DLP, NORTH WARNBORO Pit Number: 1P

Grid Reference: SU73505440 Average Annual Rainfall: 739 mm

Accumulated Temperature: 1435 degree days

Field Capacity Level : 158 days
Land Use : Arable
Slope and Aspect : degrees

| HORIZON | TEXTURE | COLOUR    | STONES >2 | TOT.STONE | LITH | MOTTLES | STRUCTURE | CONSIST | SUBSTRUCTURE | CALC |
|---------|---------|-----------|-----------|-----------|------|---------|-----------|---------|--------------|------|
| 0- 30   | MZCL    | 10YR42 43 | 2         | 8         | HR   |         |           |         |              | Y    |
| 30- 46  | MZCL    | 10YR44 00 | 0         | 5         | HR   |         | MDCSAB    | FR      | M            | Y    |
| 46- 88  | MZCL    | 10YR64 81 | 0         | 25        | HR   |         | WKCSAB    | FR      | M            | Y    |
| 88-110  | CH      | 10YR81 00 | 0         | 5         | HR   |         |           |         | Р            | Υ    |

Wetness Grade: 1 Wetness Class : I

Gleying : cm SPL : No SPL

Drought Grade: 2 APW: 124mm MBW: 19 mm

APP: 110mm MBP: 12 mm

FINAL ALC GRADE : 2

MAIN LIMITATION : Droughtiness

### SOIL PIT DESCRIPTION

Site Name: HART DLP, NORTH WARNBORO Pit Number: 2P

Grid Reference: SU73605440 Average Annual Rainfall: 739 mm

Accumulated Temperature: 1435 degree days

Field Capacity Level : 158 days
Land Use : Arable

Slope and Aspect : 03 degrees N

| HORIZON | TEXTURE | COLOUR    | STONES >2 | TOT.STONE | LITH | MOTTLES | STRUCTURE | CONSIST | SUBSTRUCTURE | CALC |
|---------|---------|-----------|-----------|-----------|------|---------|-----------|---------|--------------|------|
| 0- 26   | MZCL    | 10YR43 00 | 2         | 7         | HR   |         |           |         |              | Y    |
| 26- 33  | HZCL    | 10YR44 54 | 0         | 2         | HR   |         | MDCSAB    | FM      | М            | Y    |
| 33- 65  | С       | 05Y 58 00 | 0         | 2         | HR   | F       | MDCSAB    | FM      | M            | Y    |
| 65- 85  | СН      | 10YR81 00 | 0         | 5         | HR   |         |           |         | P            | γ    |

Wetness Grade : 1 Wetness Class : I

Orought Grade : 2 APW : 110mm MBW : 5 mm

APP: 113mm MBP: 15 mm

FINAL ALC GRADE : 2

MAIN LIMITATION : Droughtiness

program: ALC012

# LIST OF BORINGS HEADERS 30/06/95 HART DLP, NORTH WARNBORO

| SAMI | PLE        | ,     | ASPECT |       |      |     | WETI  | NESS  | -WHI | EAT- | -PC | )T\$- | M.         | REL   | EROSN | FR | OST  | CHEM  | ALC |               |
|------|------------|-------|--------|-------|------|-----|-------|-------|------|------|-----|-------|------------|-------|-------|----|------|-------|-----|---------------|
| ١٥.  | GRID REF   | USE   |        | GRONT | GLEY | SPL | CLASS | GRADE | AP   | MB   | AP  | МВ    | DRT        | FL00D | Ε     | XP | DIST | LIMIT |     | COMMENTS      |
| _ 1  | SU73505450 | ) ARA |        |       |      |     | 1     | 1     | 99   | -6   | 107 | 9     | ЗА         |       |       |    |      | DR    | 2   | Imp 62 flints |
| 11   | SU7350544  | O ARA |        |       |      |     | 1     | 1     | 124  | 19   | 110 | 12    | 2          |       |       |    |      | DR    | 2   | Chalk at 88   |
| 2    | SU73605456 | ) ARA |        |       |      |     | 1     | 1     | 100  | -5   | 109 | 11    | ЗА         |       |       |    |      | ÐR    | 2   | Imp 65 flints |
| 21   | SU73605440 | ) ARA | N      | 03    |      |     | 1     | 1     | 110  | 5    | 113 | 15    | 2          |       |       |    |      | DR    | 2   | Chalk at 65   |
| 3    | SU73505440 | ) ARA |        |       |      |     | 1     | 1     | 104  | -1   | 104 | 6     | <b>3</b> A |       |       |    |      | DR    | 2   | Imp 55 flints |
| 4    | SU73605440 | ) ARA | N      | 03    | 025  |     | 2     | 2     | 108  | 3    | 107 | 9     | 3A         |       |       |    |      | WD    | 2   | Imp 70 Chalk  |

page 1

65-90 ch

10YR81 00

----MOTTLES---- PED ----STONES---- STRUCT/ SUBS AMPLE DEPTH TEXTURE COLOUR COL ABUN CONT COL. GLEY >2 >6 LITH TOT CONSIST STR POR IMP SPL CALC 1 0 HR 0-28 mzcl 10YR43 00 3 М 28-55 10YR54 00 0 0 HR 5 mzc1 55-62 mzc1 10YR64 00 0 0 CH 15 γ Imp flints 1P 0-30 mzc1 10YR42 43 2 0 HR 8 30-46 mzc1 5 MDCSAB FR M 10YR44 00 0 0 HR Υ 46-88 mzcl 10YR64 81 0 0 HR 25 WKCSAB FR M 88-110 ch 10YR81 00 0 0 HR P V.hard from 105 0-28 mzc? 10YR43 00 2 0 HR 6 28-55 mzcl 10YR54 00 0 0 HR 5 55-65 hzc1 10YR56 00 0 0 HR 10 Υ Imp flints 0-26 mzc1 10YR43 00 2 0 HR 26-33 hzc1 10YR44 54 0 0 HR 2 MDCSAB FM M Υ 33-65 c 05Y 58 00 00MN00 00 F 0 0 HR 2 MDCSAB FM M 65-85 ch 10YR81 00 0 0 HR 5 Υ 0-28 mzc1 10YR42 43 00MN00 00 F 0 0 HR 8 28-48 hzcl 10YR44 54 O O HR 5 М 48-55 mzcl 10YR64 81 0 0 CH 50 Ρ Imp flints 0-25 mzcl 10YR43 00 2 0 HR 7 05YR58 00 75YR53 00 C 25-65 с Y 0 0 HR 10

0 0 HR

5