A1

East Hampshire Local Plan
Site 1120: Land At Farnham Road,
West Liss.
Agricultural Land Classification,
ALC Map and Report.
March 1995

### AGRICULTURAL LAND CLASSIFICATION REPORT

# EAST HAMPSHIRE LOCAL PLAN SITE 1120: LAND AT FARNHAM ROAD, WEST LISS.

#### 1. Summary

- 1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality for a number of sites in the East Hampshire District. The work forms part of MAFF's statutory input to the preparation of the East Hampshire Local Plan.
- 1.2 The site comprises 1.9 hectares of land to the south of West Liss in Hampshire. An Agricultural Land Classification (ALC) survey was carried out during February 1995. The survey was undertaken at a detailed level of approximately one boring per hectare of agricultural land surveyed. A total of 2 borings and one soil inspection pit were described in accordance with MAFF's revised guidelines and criteria for grading the quality of agricultural land (MAFF, 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose a long term limitation on its use for agriculture.
- 1.4 At the time of the survey the agricultural land was under stubble from a previous cereal crop, being part of a larger field extending southwards. The area of Non-agricultural land comprises the garden to houses known as 'The Brows' immediately to the south of the site. These are separated from the agricultural land by a metalled track, shown as Urban, leading to agricultural buildings and light industry also to the south of the site.
- 1.5 The distribution of grades and subgrades is shown on the attached ALC map and the areas and extent are given in the table below. The map has been drawn at a scale of 1:10,000. It is accurate at this scale, but any enlargement would be misleading.

**Table 1: Distribution of Grades and Subgrades** 

Grade	Area (ha)	% of Site
3b	1.5	78.9
Urban	0.1	5.3
Non-Agricultural	<u>0.3</u>	<u>15.8</u>
Total area of site	1.9ha	100.0

1.6 Appendix I gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.

1.7 The agricultural land at this site has been classified as moderate quality (Subgrade 3b). The principal limitation to land quality is soil wetness due to poorly structured slowly permeable clay subsoils significantly impeding drainage.

#### 2. Climate

- 2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.
- 2.2 The main parameters used in the assessment of an overall climatic limitation are average annual rainfall, as a measure of overall wetness, and accumulated temperature, as a measure of the relative warmth of a locality.
- A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met. Office, 1989). The details are given in the table below and these show that there is no overall climatic limitation affecting the site.
- 2.4 No local climatic factors such as exposure or frost risk are believed to affect the site. However, climatic and soil factors interact to influence soil wetness and droughtiness limitations.

**Table 2: Climatic Interpolation** 

Grid Reference	SU774280
Altitude, (m, AOD)	70
Accumulated Temperature	1462
(°days, JanJune)	
Average Annual Rainfall (mm)	905
Field Capacity Days	200
Moisture deficit, wheat (mm)	97
Moisture deficit, potatoes (mm)	88
Overall Climatic Grade	l

#### 3. Relief

3.1 The site lies at approximately 70m AOD. Overall the site falls slightly from north west to south east. Nowhere in this area does relief or gradient affect agricultural land quality.

### 4. Geology and Soils

- 4.1 The published geological information (BGS, 1981), shows the site to be underlain by Cretaceous Folkestone Beds.
- 4.2 The published soils information (SSEW, 1983), shows the site to be underlain by soils of the Fyfield 4 Association. The legend accompanying the map describes these as, 'deep well drained often stoneless coarse loamy and sandy soils. Some fine loamy soils with slowly permeable subsoils and slight seasonal waterlogging

and some slowly permeable seasonally waterlogged fine loamy over clayey soils. Risk of water erosion.' (SSEW, 1983). Soils encountered at this site comprised fine loamy topsoils over slowly permeable clayey subsoils.

### 5. Agricultural Land Classification

- 5.1 Paragraph 1.5 provides the details of the area measurements for each grade and the distribution of each grade is shown on the attached ALC map.
- 5.2 The location of the soil observation points are shown on the attached sample point map.

### Subgrade 3b

5.3 Land of moderate quality has been mapped over the agricultural area of this site. The principal limitation is soil wetness, due to poor drainage. Soil profiles typically comprise a stoneless or very slightly stony (0-3% total flints), occasionally gleyed medium sandy loam or sandy clay loam topsoil. This either passes directly to a stoneless, gleyed and slowly permeable clay (see Pit 1, Appendix III), or has a gleyed heavy clay loam upper subsoil overlying the clay. The clay severely impedes drainage and soils are assigned to Wetness Class IV. The soil wetness which occurs will adversely affect crop growth and development and restricts opportunities for landwork and/or grazing by livestock. Occasionally the clay is overlain by gleyed medium sandy loam and sandy clay loam horizons creating a profile of slightly better quality.

ADAS Ref: 1502/012/95 MAFF Ref: EL15/468

Resource Planning Team Guildford Statutory Group ADAS Reading

### **SOURCES OF REFERENCE**

British Geological Survey (1975), Sheet 300, Alresford, Drift Edition. 1:50,000

MAFF (1988), Agricultural Land Classification of England and Wales: Revised guidelines and criteria for grading the quality of agricultural land.

Meteorological Office (1989), Climatic datasets for Agricultural Land Classification.

Soil Survey of England and Wales (1983), Sheet No. 6, Soils of South-East England, 1:250,000, and Accompanying Legend.

Soil Survey of England and Wales (1984), Bulletin No.15, Soils and their use in South-East England.

### APPENDIX I

#### DESCRIPTION OF THE GRADES AND SUBGRADES

### Grade 1: Excellent Quality Agricultural Land

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

# Grade 2: Very Good Quality Agricultural Land

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land of this grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1 land.

### Grade 3: Good to Moderate Quality Land

Land with moderate limitations which affect the choice of crops, the timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown, yields are generally lower or more variable than on land in Grades 1 and 2.

### Subgrade 3a: Good Quality Agricultural Land

Land capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

### Subgrade 3b: Moderate Quality Agricultural Land

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

### Grade 4: Poor Quality Agricultural Land

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (e.g. cereals and forage crops) the yields of which are variable. In moist climates, yields of grass may be moderate to high but there may be difficulties in utilisation. The grade also includes very droughty arable land.

### Grade 5: Very Poor Quality Agricultural Land

Land with severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

#### Urban

Built-up or 'hard' uses with relatively little potential for a return to agriculture including: housing, industry, commerce, education, transport, religious buildings, cemeteries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

### Non-agricultural

'Soft' uses where most of the land could be returned relatively easily to agriculture, including: private parkland, public open spaces, sports fields, allotments and soft-surfaced areas on airports. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

#### Woodland

Includes commercial and non-commercial woodland. A distinction may be made as necessary between farm and non-farm woodland.

#### **Agricultural Buildings**

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (e.g. polythene tunnels erected for lambing) may be ignored.

#### Open Water

Includes lakes, ponds and rivers as map scale permits.

#### Land Not Surveyed

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above, e.g. buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will be shown.

### APPENDIX II

### **DEFINITION OF SOIL WETNESS CLASS**

### Wetness Class I

The soil profile is not wet within 70 cm depth for more than 30 days in most years.

#### Wetness Class II

The soil profile is wet within 70 cm depth for 31-90 days in most years or, if there is no slowly permeable layer within 80 cm depth, it is wet within 70 cm for more than 180 days, but only wet within 40 cm depth for 31-90 days in most years.

#### Wetness Class III

The soil profile is wet within 70 cm depth for 91-180 days in most years or, if there is no slowly permeable layer present within 80 cm depth, it is wet within 70 cm for more than 180 days, but only wet within 40 cm depth for between 31-90 days in most years.

### Wetness Class IV

The soil profile is wet within 70 cm depth for more than 180 days but not wet within 40 cm depth fro more than 210 days in most years or, if there is no slowly permeable layer present within 80 cm depth, it is wet within 40 cm depth for 91-210 days in most years.

#### Wetness Class V

The soil profile is wet within 40 cm depth for 211-335 days in most years.

#### Wetness Class VI

The soil profile is wet within 40 cm depth for more than 335 days in most years.

# APPENDIX III

# SOIL PIT AND SOIL BORING DESCRIPTIONS

### Contents:

Sample Point Map

Soil Abbreviations - explanatory note

Database Printout - soil pit information

**Database Printout - boring level information** 

**Database Printout - horizon level information** 

## SOIL PROFILE DESCRIPTIONS: EXPLANATORY NOTE

Soil pit and auger boring information collected during ALC fieldwork is held on a database. This has commonly used notations and abbreviations as set out below.

### **Boring Header Information**

1. **GRID REF**: national grid square and 8 figure grid reference.

2. USE: Land use at the time of survey. The following abbreviations are used.

ARA: Arable WHT: Wheat BAR: Barley
CER: Cereals OAT: Oats MZE: Maize
OSR: Oilseed rape BEN: Field Beans BRA: Brassicae
POT: Potatoes SBT: Sugar Beet FCD: Fodder Crops

LIN: Linseed FRT: Soft and Top Fruit FLW: Fallow

PGR: Permanent Pasture LEY: Ley Grass RGR: Rough Grazing SCR: Scrub CFW: Coniferous Woodland DCW: Deciduous Wood

HTH: Heathland BOG: Bog or Marsh FLW: Fallow PLO: Ploughed SAS: Set aside OTH: Other

**HRT**: Horticultural Crops

3. **GRDNT**: Gradient as measured by a hand-held optical clinometer.

4. **GLEY/SPL**: Depth in cm to gleying or slowly permeable layers.

5. **AP (WHEAT/POTS)**: Crop-adjusted available water capacity.

6. **MB (WHEAT/POTS)**: Moisture Balance.

7. **DRT**: Best grade according to soil droughtiness.

8. If any of the following factors are considered significant, an entry of 'Y' will be entered in the relevant column.

MREL: Microrelief limitation FLOOD: Flood risk EROSN: Soil erosion risk EXP: Exposure limitation FROST: Frost DIST: Disturbed land

**CHEM**: Chemical limitation

9. **LIMIT**: The main limitation to land quality. The following abbreviations are used.

FL: Flood Risk TX: Topsoil Texture DP: Soil Depth ST: Topsoil Stones

CH: Chemical WE: Wetness WK: Workability

**DR**: Drought **ER**: Erosion Risk **WD**: Soil Wetness/Droughtiness

# Soil Pits and Auger Borings

1. **TEXTURE**: soil texture classes are denoted by the following abbreviations.

S: Sand LS: Loamy Sand SL: Sandy Loam

SZL: Sandy Silt Loam CL: Clay Loam

ZCL: Silty Clay Loam SCL: Sandy Clay Loam

C: Clay SC: Sandy Clay Clay SP: Sandy Peat PE: Peaty Loam PS: Peaty Sand

MZ: Marine Light Silts

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction will be indicated by the use of prefixes.

**F**: Fine (more than 66% of the sand less than 0.2mm)

M: Medium (less than 66% fine sand and less than 33% coarse sand)

C: Coarse (more than 33% of the sand larger than 0.6mm)

The clay loam and silty clay loam classes will be sub-divided according to the clay content: M : Medium (<27% clay) H : Heavy (27-35% clay)

- 2. MOTTLE COL: Mottle colour
- 3. MOTTLE ABUN: Mottle abundance, expressed as a percentage of the matrix or surface described.

F: few <2% C: common 2-20% M: many 20-40 VM: very many 40%

4. MOTTLE CONT : Mottle contrast

**F**: faint - indistinct mottles, evident only on close inspection

**D**: distinct - mottles are readily seen

P: prominent - mottling is conspicuous and one of the outstanding features of the horizon

- 5. **PED. COL**: Ped face colour
- 6. **STONE LITH**: One of the following is used.

HR: all hard rocks and stones SLST: soft oolitic or dolimitic limestone

CH: chalk FSST: soft, fine grained sandstone

**ZR**: soft, argillaceous, or silty rocks **GH**: gravel with non-porous (hard) stones **MSST**: soft, medium grained sandstone **GH**: gravel with non-porous (hard) stones

**S1**: soft weathered igneous/metamorphic rock

Stone contents (>2cm, >6cm and total) are given in percentages (by volume).

7. STRUCT: the degree of development, size and shape of soil peds are described using the following notation:

degree of development WK: weakly developed MD: moderately developed

**ST**: strongly developed

ped size ped shape
 F: fine M: medium C: coarse VC: very coarse
 S: single grain M: massive GR: granular AB: angular blocky

**SAB**: sub-angular blocky **PR**: prismatic **PL**: platy

8. **CONSIST**: Soil consistence is described using the following notation:

L: loose VF: very friable FR: friable FM: firm VM: very firm EM:

extremely firm

EH: extremely hard

9. SUBS STR: Subsoil structural condition recorded for the purpose of calculating profile droughtiness: G: good M: moderate P: poor

- 10. POR: Soil porosity. If a soil horizon has less than 0.5% biopores >0.5 mm, a 'Y' will appear in this column.
- 11. **IMP**: If the profile is impenetrable a 'Y' will appear in this column at the appropriate horizon.
- 12. SPL: Slowly permeable layer. If the soil horizon is slowly permeable a 'Y' will appear in this column.
- 13. CALC: If the soil horizon is calcareous, a 'Y' will appear in this column.
- 14. Other notations

**APW**: available water capacity (in mm) adjusted for wheat **APP**: available water capacity (in mm) adjusted for potatoes

MBW: moisture balance, wheat MBP: moisture balance, potatoes

SOIL PIT DESCRIPTION

Site Name : E HANTS LP SITE 1120 Pit Number : 1P

Grid Reference: SU77392796 Average Annual Rainfall: 905 mm

Accumulated Temperature : 1462 degree days

Field Capacity Level : 200 days

Land Use

Slope and Aspect : degrees

HORIZON	TEXTURE	COLOUR	STONES >2	TOT.STONE	LITH	MOTTLES	STRUCTURE	CONSIST	SUBSTRUCTURE	CALC
0- 34	SCL	10YR42 00	0	3	HR					
34- 56	С	05Y 62 63	0	0	HR	С	WKCSAB	FM	M	
56- 80	С	05GY71 00	0	0	HR	С	WKCSAB	FM	М	
80- 90	С	05GY51 00	0	0	HR.	С			Р	
90-110	SCL	05GY61 00	0	0	HR	С			М	
110-120	С	05GY51 00	0	ó	HR	С			₽	

:

Wetness Grade : 3B Wetness Class : IV

Gleying : 34 cm SPL : 34 cm

Drought Grade: APW: mm MBW: 0 mm

APP: mm MBP: 0 mm

FINAL ALC GRADE : 3B
MAIN LIMITATION : Wetness

program: ALCO12

# LIST OF BORINGS HEADERS 28/02/95 E HANTS LP SITE 1120

page 1

.

	SAMP	LE	A	SPECT				WETN	VESS	-WHE	AT-	-P0	TS-	м. я	REL	EROSN	FROST	CHEM	ALC	
ì	NO.	GRID REF	USE		GRDNT	GLEY	SPL	CLASS	GRADE	ΑP	MB	AP	MB	DRT	FLOOD	EX	KP DIST	LIMIT		COMMENTS
	1	SU77352800	STB			30	75	3	3A		0		D					WE	3A	
_	1P	SU77392796	STB			34	34	4	38		0		0					WE	3В	PIT 80 AUG 120
ı	2	SU77442793	STB	SE	01	Q	50	4	3B		0		0					WE	38	

				M	OTTL	ES	PED			·-S	TONES-		STRUCT/	' SUB	\$			
SAMPLE	DEPTH	TEXTURE	COLOUR	COL	NUBA	CONT	COL.	GLEY	>2	>6	LITH	TOT	CONSIST	STR	POR	IMP	SPL	CALC
1	0-30	msl	10YR42 00						0	0	HR	0						
	30-50	ms 1	25Y 62 63	10YR66	00	С		Υ	0	0	HR	0		M				
	50-75	foa	25Y 62 00	10YR58	00	C		Υ	0	0	HR	0		М				
	75-120	С	25Y 61 62	10YR58	00	С		Υ	0	0	HR	0		Р			Y	
		,	400040 00							_		_						
1P	0-34	scl	10YR42 00						0	U	HR	3						
	34-56	С	05Y 62 63	10YR58	00	С		Υ	0	0	HR	0	WKCSAB	FM M	Υ		Y	
	56-80	С	05GY71 00	10YR68	00	С		Υ	0	0	HR	0	WKCSAB	FM M	Y		Υ	
	80-90	С	05GY51 00	10YR68	00	С		Υ	0	0	HR	0		Р			Υ	
	90-110	scl	05GY61 00	10YR66	00	С		Υ	0	0	HR	0		М			γ	
	110-120	С	05GY51 00	10YR68	00	С	i	Υ	0	0	HR	0		Р			Υ	
2	0-30	scl	25Y 41 00	10YR46	00	c ·		Υ	0	0	HR	2						
	30-50	hc1	25Y 61 00	10YR68	00	М		Υ	0	0		0		М				
	50-90	С	05Y 61 00	10YR68	00	М		Υ	0	0		0		Р			Υ	