A1
Newbury District Local Plan
Site 48: Greenham Road, Greenham
Agricultural Land Classification
ALC Map and Report
February 1994

NEWBURY DISTRICT LOCAL PLAN SITE 48: GREENHAM ROAD, GREENHAM, BERKSHIRE AGRICULTURAL LAND CLASSIFICATION REPORT

1. Summary

- 1.1 ADAS was commissioned by MAFF's Land Use Planning Unit to provide information on land quality for a number of sites in the Newbury District of Berkshire. The work forms part of MAFF's statutory input to the preparation of the Newbury District Local Plan.
- 1.2 Approximately 20 hectares of land relating to site 48, Greenham Road, Greenham, Berkshire was surveyed in February 1994. The survey was undertaken at a detailed level of approximately one observation per hectare. A total of 15 soil auger borings and three soil inspection pits were assessed in accordance with MAFF's revised guidelines and criteria for grading the quality of agricultural land (MAFF, 1988). These guidelines provide a framework for classifying land according to the extent to which its physical or chemical characteristics impose long term limitations on its use for agriculture.
- 1.3 The work was conducted by members of the Resource Planning Team in the Guildford Statutory Group of ADAS.
- 1.4 At the time of the survey the land under agricultural use was in grass, towards the north east of the site this was unmanaged and overgrown. The remainder being permanent grazing or a ley.
- 1.5 The distribution of grades and subgrades is shown on the attached ALC map and the areas and extent are given in the table below. The map has been drawn at a scale of 1:5,000. It is accurate at this scale, but any enlargement would be misleading. This map supersedes a previous survey carried out in 1988, by ADAS, for the proposed Newbury by-pass.

Table 1: Distribution of Grades and Subgrades

Grade	Area(ha)	% of Site	% of Agricultural Area
3b	9.5	48.2	76.0
4	3.0	15.2	<u>24.0</u>
Non-agricultural	0.4	2.0	100% (12.5 ha)
Woodland	5,5	28.0	
Urban	1.2	6.1	
Agricultural Buildings	<u>0.1</u>	<u>0.5</u>	
Total Area of Site	19.7 ha	100%	

1.6 Appendix 1 gives a general description of the grades, subgrades and land use categories identified in the survey. The main classes are described in terms of the type of limitation that can occur, the typical cropping range and the expected level and consistency of yield.

1.7 The land has been classified as Subgrade 3b (moderate quality) and Grade 4 (poor quality). Limitations include soil droughtiness, soil wetness and gradient. Towards the north of the site soil wetness is the principal limitation; soils here are characterised by moderately stony shallow slowly permeable clay horizons. These poorly drained soils are appropriately placed in Subgrade 3b. Towards the south of the site, on higher land, soil droughtiness becomes the principal limitation. This is due to moderately and extremely stony soil profiles restricting water availability to plants such that there is a risk of significant or severe drought stress. Moisture balances indicate Subgrade 3b and Grade 4 are appropriate given the local climatic conditions. The small area shown as Grade 4, towards the east of the site is limited by a gradient of 13 degrees. This affects the safe and efficient use of farm machinery such that regular cultivation may be difficult.

2. Climate

- 2.1 The climatic criteria are considered first when classifying land as climate can be overriding in the sense that severe limitations will restrict land to low grades irrespective of favourable site or soil conditions.
- 2.2 The main parameters used in the assessment of an overall climatic limitation are average annual rainfall, as a measure of overall wetness, and accumulated temperature, as a measure of the relative warmth of a locality.
- 2.3 A detailed assessment of the prevailing climate was made by interpolation from a 5km gridpoint dataset (Met. Office, 1989). The details are given in the table below and these show that there is no overall climatic limitation affecting the site.
- 2.4 No local climatic factors such as exposure or frost risk affect the site. However, climatic and soil factors interact to influence soil wetness and droughtiness limitations.

Table 2: Climatic Interpolation

Grid Reference:	SU 484660	SU484655
Altitude (m):	90	120
Accumulated Temperature (°days):	1429	1395
Average Annual Rainfall (mm):	726	754
Field Capacity (days):	159	163
Moisture Deficit, Wheat (mm):	105	100
Moisture Deficit, Potatoes (mm):	96	91
Overall Climatic Grade:	1	1

3. Relief

3.1 The site lies at approximately 90 m and 120 m AOD, rising from north to south.

Relief and microrelief do not generally affect the land quality on the site, however in a small area east of West Wood gradients of c.13° affects the safe and efficient use of farm machinery to the extent that Grade 4 is appropriate.

4. Geology and Soil

- 4.1 The published British Geological Survey map, Sheet 267, Hungerford (1971, 1:63,360 scale), shows the north of the site to be underlain by London Clay. To the south of the site plateau gravel is mapped, where the land is higher.
- 4.2 The published Soil Survey of England and Wales map, Sheet 6, Soils of South East England (1983, 1:250,000 scale), shows the site as the Sonning 2 Association. These soils are described by the SSEW as "well drained flinty coarse loamy and gravelly soils. Associated with slowly permeable seasonally waterlogged fine loamy over clayey soils, and coarse loamy over clayey soils with slowly permeable subsoils and slight seasonal waterlogging. (SSEW, 1983). Soils of this broad nature were encountered across the site during detailed ALC survey work; towards the north east slowly permeable subsoils were found and towards the south and west the profiles were found to be gravelly.

5. Agricultural Land Classification

- Table 1 provides the details of the area measurements for each grade and the distribution of each grade is shown on the attached ALC map.
- 5.2 The location of the soil observation points are shown on the attached sample point map.

Subgrade 3b

5.3 Land of moderate quality occurs across the north and west of the site in a single mapping unit. Soils in this area were found to be of two distinct types. Towards the north east of the site soil wetness was the principal limitation. In this area soils were characterised by a very slightly stony medium clay loam topsoil, which either overlies a moderately stony heavy clay loam passing to a stoneless or moderately to very stony slowly permeable clay, or, passes directly to the clay. These soils are poorly drained (Wetness Class IV, see Appendix II), and are therefore limited to Subgrade 3b on this basis. Towards the south and west of the site soil droughtiness becomes the principal limitation. In this area soils are characterised by a moderately stony (max 17% v/v flints > 2 cm) medium clay loam topsoil, which passes to a very stony (c.60% v/v flints) fine loamy or coarse loamy upper subsoil becoming gravel within 50 cm. These soils are well drained (Wetness Class I, see Appendix II), but are limited to Subgrade 3b by soil droughtiness caused by the presence of flints in the soil profile often in combination with topsoil stone contents which are also limiting. Stones reduce the available water capacity in the profile such that there is a

significant risk of drought stress to crops. This results in lower and/or less consistent yields of a narrow range of crops, principally cereals and grass.

Grade 4

- Land of poor quality is mapped towards the south-east and east of the site in two mapping units. The principal limitation is severe soil droughtiness, due primarily to high stone contents restricting water availability to crops. Soils typically comprise a slightly stony (c.15% v/v flints) medium clay loam topsoil, passing to a shallow (c.5 cm) very stony (c.50% v/v flints) medium clay loam upper subsoil overlying gravel at c.20-28 cm. Soils are stonier than for similar land graded 3b, such that crops are at a greater risk of drought stress. A small area towards the east of West Wood is limited by gradient, measured using an optical reading clinometer, at 13°. This affects the safe and efficient use of farm machinery such that regular cultivation my be difficult. Land of this quality could be expected to produce very variable yields of a narrow range of crops, principally grass, and occasional arable crops such as cereals or forage crops.
- 5.5 The areas shown as Urban at this site consist of domestic dwellings, associated grounds and driveways, and a fenced metalled track (New Road). The areas shown as non-agricultural are Hawthorn scrub. The areas shown as agricultural buildings are battery hen houses which are now derelict, the areas between them being overgrown and shown as non-agricultural.

ADAS Ref: 0202/011/94 MAFF Ref: EL02/0297 Resource Planning Team Guildford Statutory Group ADAS Reading

SOURCES OF REFERENCE

- * British Geological Survey (1971), Sheet No. 267, Hungerford, 1:63,360.
- * MAFF (1988), Agricultural Land Classification of England and Wales. Revised guidelines and criteria for grading the quality of agricultural land.
- * Meteorological Office (1989), Climatological Data for Agricultural Land Classification.
- * Soil Survey of England and Wales (1983), Sheet No 6, Soils of South East England, 1:250,000, and Accompanying Legend.
- * Soil Survey of England and Wales (1984), Soils and their use in South East England. Bulletin No. 15.

APPENDIX I

DESCRIPTION OF THE GRADES AND SUBGRADES

Grade 1: Excellent Quality Agricultural Land

Land with no or very minor limitations to agricultural use. A very wide range of agricultural and horticultural crops can be grown and commonly includes top fruit, soft fruit, salad crops and winter harvested vegetables. Yields are high and less variable than on land of lower quality.

Grade 2: Very Good Quality Agricultural Land

Land with minor limitations which affect crop yield, cultivations or harvesting. A wide range of agricultural or horticultural crops can usually be grown but on some land of this grade there may be reduced flexibility due to difficulties with the production of the more demanding crops such as winter harvested vegetables and arable root crops. The level of yield is generally high but may be lower or more variable than Grade 1 land.

Grade 3: Good to Moderate Quality Land

Land with moderate limitations which affect the choice of crops, the timing and type of cultivation, harvesting or the level of yield. When more demanding crops are grown, yields are generally lower or more variable than on land in Grades 1 and 2.

Subgrade 3a: Good Quality Agricultural Land

Lánd capable of consistently producing moderate to high yields of a narrow range of arable crops, especially cereals, or moderate yields of a wide range of crops including cereals, grass, oilseed rape, potatoes, sugar beet and the less demanding horticultural crops.

Subgrade 3b: Moderate Quality Agricultural Land

Land capable of producing moderate yields of a narrow range of crops, principally cereals and grass, or lower yields of a wider range of crops or high yields of grass which can be grazed or harvested over most of the year.

Grade 4: Poor Quality Agricultural Land

Land with severe limitations which significantly restrict the range of crops and/or the level of yields. It is mainly suited to grass with occasional arable crops (eg. cereals and forage crops) the yields of which are variable. In moist climates, yields of grass may be moderate to high but there may be difficulties in utilisation. The grade also includes very droughty arable land.

Grade 5: Very Poor Quality Agricultural Land

Land with severe limitations which restrict use to permanent pasture or rough grazing, except for occasional pioneer forage crops.

Urban

Built-up or 'hard' uses with relatively little potential for a return to agriculture including: housing, industry, commerce, education, transport, religous buildings, cemetries. Also, hard-surfaced sports facilities, permanent caravan sites and vacant land; all types of derelict land, including mineral workings which are only likely to be reclaimed using derelict land grants.

Non-agricultural

'Soft' uses where most of the land could be returned relatively easily to agriculture, including: private parkland, public open spaces, sports fields, allotments and soft-surfaced areas on airports. Also active mineral workings and refuse tips where restoration conditions to 'soft' after-uses may apply.

Woodland

Includes commercial and non-commercial woodland. A distinction may be made as necessary between farm and non-farm woodland.

Agricultural Buildings

Includes the normal range of agricultural buildings as well as other relatively permanent structures such as glasshouses. Temporary structures (eg. polythene tunnels erected for lambing) may be ignored.

Open Water

Includes lakes, ponds and rivers as map scale permits.

Land Not Surveyed

Agricultural land which has not been surveyed.

Where the land use includes more than one of the above, eg. buildings in large grounds, and where map scale permits, the cover types may be shown separately. Otherwise, the most extensive cover type will be shown.

APPENDIX II

FIELD ASSESSMENT OF SOIL WETNESS CLASS

Definition of Soil Wetness Classes

Wetness Class	Duration of Waterlogging ¹
I	The soil profile is not wet within 70 cm depth for more than 30 days in most years ² .
II	The soil profile is wet within 70 cm depth for 31-90 days in most years <u>or</u> , if there is no slowly permeable layer within 80 cm depth, it is wet within 70 cm for more than 90 days, but not wet within 40 cm depth for more than 30 days in most years.
Ш	The soil profile is wet within 70 cm depth for 91-180 days in most years <u>or</u> , if there is no slowly permeable layer within 80 cm depth, it is wet within 70 cm for more than 180 days, but only wet within 40 cm depth for between 31 and 90 days in most years.
· IV	The soil profile is wet within 70 cm depth for more than 180 days but not within 40 cm depth for more than 210 days in most years <u>or</u> , if there is no slowly permeable layer within 80 cm depth, it is wet within 40 cm depth for 91-210 days in most years.
V	The soil profile is wet within 40 cm depth for 211-335 days in most years.
VI	The soil profile is wet within 40 cm depth for more than 335 days in most years.

¹ The number of days specified is not necessarily a continuous period.

² 'In most years' is defined as more than 10 out of 20 years.

APPENDIX III

SOIL PIT AND SOIL BORING DESCRIPTIONS

Contents:

Soil Abbreviations - Explanatory Note

Soil Pit Descriptions

Database Printout - Boring Level Information

Database Printout - Horizon Level Information

SOIL PROFILE DESCRIPTIONS: EXPLANATORY NOTE

Soil profile and pit information obtained during ALC surveys is held on a database. This has commonly used notations and abbreviations as set out below.

BORING HEADERS

- 1. GRID REF: National grid square followed by 8 figure grid reference.
- 2. USE: Land-use at the time of survey.

 The following abbreviations are used.

ARA - arable PAS/PGR - permanent pasture WHT - wheat RGR - rough grazing BAR - barley LEY - ley grassland CER - cereals CFW - coniferous woodland OAT - oats DCW - deciduous woodland SCR - scrub MZE - maize OSR - oilseed rape HTH - heathland BEN - field beans BOG - bog or marsh BRA - brassicae FLW - fallow POT - potatoes PLO - ploughed SBT - sugarbeet SAS - set-aside FDC - fodder crops OTH - other FRT - soft and top fruit LIN - linseed HOR/HRT - horticultural crops

- 3. GRDNT: Gradient as measured by optical reading clinometer.
- 4. GLEY/SPL: Depth in centimetres (cm) to gleyed and/or slowly permeable horizons.
- 5. AP (WHEAT/POTS) : Crop-adjusted available water capacity. The amount of soil water (in millimetres) held in the soil profile that is available to a growing crop (wheat and potatoes are used as reference crops).
- 6. MB (WHEAT/POTS): The moisture balance for wheat and potatoes obtained by subtracting the soil moisture deficit from the crop-adjusted available water capacity.
- 7. DRT: Grade according to soil droughtiness assessed against soil moisture balances.

```
8. M REL: Micro-relief
FLOOD: Flood risk
EROSN: Soil erosion
EXP: Exposure
FROST: Frost prone
DIST: Disturbed land
CHEM: Chemical limitation)

If any of these factors are considered significant in terms of the assessment of agricultural land quality a 'y' will be entered in the relevant column.
```

9. LIMIT: Principal limitation to agricultural land quality.
The following abbreviations are used:

OC - overall climate

AE - aspect

EX - exposure

FR - frost

GR - gradient

CH - chemical limitations

WE - wetness

WK - workability

DR - drought

ER - erosion

MR- micro-relief WD - combined soil wetness/soil
FL - flooding droughtiness
TX - soil texture ST - topsoil stoniness

DP - soil depth

PROFILES & PITS

1. TEXTURE: Soil texture classes are denoted by the following abbreviations:

S - sand '

LS - loamy sand
SL - sandy loam
SZL - sandy silt loam

ZL - silt loam

MZCL - medium silty clay loam

MCL - medium clay loam

SCL - sandy clay loam

HZCL - heavy silty clay loam

HCL - heavy clay loam

SC - sandy clay
ZC - silty clay

C - clay

For the sand, loamy sand, sandy loam and sandy silt loam classes, the predominant size of sand fraction may be indicated by the use of prefixes.

F - fine (more than ²/₃ of the sand less than 0.2 mm)

C - coarse (more than 1/3 of sand greater than 0.6 mm)

M - medium (less than ²/₃ fine sand and less than ¹/₃ coarse sand)

The sub-divisions of clay loam and silty clay loam classes according to clay content are indicated as follows:

M - medium (less than 27% clay)

H - heavy (27-35% clay)

Other possible texture classes include:

OL - organic loam

P - peat

SP - sandy peat

LP - loamy peat

PL - peaty loam

PS - peaty sand

MZ - marine light silts

- 2. MOTTLE COL: Mottle colour
- 3. MOTTLE ABUN: Mottle abundance

F - few - less than 2% of matrix or surface described

C - common - 2-20% of the matrix

M - many - 20-40% of the matrix

VM - very many - 40% + of the matrix

- 4. MOTTLE CONT: Mottle continuity
 - F faint indistinct mottles, evident only on close examination
 - D distinct mottles are readily seen
 - P prominent mottling is conspicuous and one of the outstanding features of the horizon
- 5. PED.COL: Ped face colour
- 6. STONE LITH: Stone lithology. One of the following is used.

HR - all hard rocks or stones

MSST - soft, medium or coarse grained sandstone

SI - soft weathered igneous or metamorphic

SLST - soft oolitic or dolomitic limestone

FSST - soft, fine grained sandstone

ZR - soft, argillaceous, or silty rocks

CH - chalk

GH - gravel with non-porous (hard) stones

GS - gravel with porous (soft) stones

Stone contents (>2cm, >6cm and total) are given in percentages (by volume).

- 7. STRUCT: the degree of development, size and shape of soil peds are described using the following notation.
 - <u>degree of development</u> WK weakly developed

MD - moderately developed

ST - strongly well developed

- ped size

F - fine

M - medium

C - coarse

VC - very coarse

- ped shape

1

S - single grain M - massive

GR - granular

SB/SAB - sub-angular blocky

AB - angular blocky

PR - prismatic

PL - platy

8. CONSIST: Soil consistence is decribed using the following notation:

L - loose

VF - very friable

FR - friable

FM - firm

VM - very firm

EM - extremely firm

EH - extremely hard

9. SUBS STR: Subsoil structural condition recorded for the purpose of calculating profile droughtiness.

G - good

M - moderate

P - poor

- 10. POR: Soil porosity. If a soil horizon has less than 0.5% biopores >0.5 mm, a 'y' will appear in this column.
- 11. IMP: If the profile is impenetrable a 'y' will appear in this column at the appropriate horizon.
- 12. SPL: Slowly permeable layer. If the soil horizon is slowly permeable a 'y' will appear in this column.
- 13. CALC: If the soil horizon is calcareous, a 'y' will appear in this column.
- 14. Other Notations

APW - available water capacity (in mm) adjusted for wheat

APP - available water capacity (in mm) adjusted for potatoes

MBW - moisture balance, wheat

MBP - moisture balance, potatoes

SOIL PIT DESCRIPTION

Site Name: NEWBURY LP SITE 48

Pit Number: 2P

Grid Reference: SU48496567

Average Annual Rainfall: 754 mm

Accumulated Temperature: 1395 degree days

Field Capacity Level : 163 days
Land Use : Rough Grazing
Slope and Aspect : 04 degrees N

HORIZON	TEXTURE	COLOUR	STONES >2	TOT.STONE	MOTTLES	STRUCTURE
0- 15	MCL	10YR32 42	0	2		
15- 28	∫HCL	10YR42 00	0	20		WKCSAB
28- 43	` c	10YR52 00	0	0	М	MDCSAB
43- 60	С	10YR53 00	0	0	М	MDCAB

Wetness Grade : 3B

Wetness Class : IV
Gleying :028 cm
SPL :043 cm

Drought Grade :

APW: mm MBW: 0 mm APP: mm MBP: 0 mm

FINAL ALC GRADE : 3B
MAIN LIMITATION : Wetness

SOIL PIT DESCRIPTION

Site Name: NEWBURY LP SITE 48

Pit Number: 3P

Grid Reference: SU48156569

Average Annual Rainfall: 754 mm

Accumulated Temperature: 1395 degree days

Field Capacity Level : 163 days Land Use : Ley

Slope and Aspect : 02 degrees N

HORIZON	TEXTURE	COLOUR	STONES >2	TOT.STONE	MOTTLES	STRUCTURE
0- 24	MCL	10YR32 42	17	41		
24- 50	CSL	10YR42 00	16	57		
50-120	GH	75YR43 00	0	0		

Wetness Grade : 1

Wetness Class : I
Gleying : cm
SPL : No SPL

Drought Grade : 3B

APW: 53 mm MBW: -47 mm APP: 50 mm MBP: -41 mm

FINAL ALC GRADE : 3B

MAIN LIMITATION : Droughtiness

SOIL PIT DESCRIPTION

Site Name: NEWBURY LP SITE 48 Pit Number: 4P

Grid Reference: SU48516564 Average Annual Rainfall: 754 mm

Accumulated Temperature: 1395 degree days

Field Capacity Level : 163 days
Land Use : Rough Grazing
Slope and Aspect : 02 degrees N

HORIZON	TEXTURE	COLOUR	STONES >2	TOT.STONE	MOTTLES	STRUCTURE
0- 15	MCL	10YR32 42	11	29		
15- 58	MCL	10YR42 00	0	55		
58- 76	MCL	10YR53 00	0	10	С	MDCSAB
76- 95	С	10YR52 00	0	5	М	
95-110	С	75YR58 00	0	30	F	

Wetness Grade : 2 Wetness Class : II

Gleying :058 cm SPL :076 cm

Drought Grade : 3A APW : 88 mm MBW : -12 mm

APP: 70 mm MBP: -21 mm

FINAL ALC GRADE : 3A

MAIN LIMITATION : Droughtiness

program: ALC012 LIST OF BORINGS HEADERS 23/03/94 NEWBURY LP SITE 48

page 1

SAMP	LE	A	SPECT				WETN	NESS	-WH	EAT-	-P0	TS-	M. F	REL	EROSN	EROSN FROST		CHEM	ALC	
NO.	GRID REF	USE		GRDNT	GLEY	SPL	CLASS	GRADE	AP	MB	AP	MB	DRT	FL00D	E	ΧP	DIST	LIMIT		COMMENTS
2P	SU48496567	RGR	N	04	028 (043	4	3B		0		0						ME	3В	PIT 60 AUG 120
3P	SU48156569	Γ₽Λ	N	02			1	1	53	-47	50	-41	3B					DR	3B	DR TO 120
30	SU48156569	LEY	N	02			1	1	47	-53	48	-43	4					DR	4	PIT59 DR TO 59
- 4	SU48426591	RGR			022		4	3B	67	-33	67	-24	3B					WD	3B	IMPST 50
4P	SU48516564	RGR	N	02	058 (076	2	2	88	-12	70	-21	3A					DR	ЗА	PIT 76 AUG 110
5	SU48506590	RGR	N	01			1	1	55	-45	55	-36	3B					DR	3B	IMPGH 37
6	SU48206579	LEY	N	06			1	1	43	-57	43	-48	4					DR	4	IMPGH 30
6 Q	SU48206579	LEY	N	06			1	1	58	-42	55	-36	3B					DR	3B	DR TO 120
9	SU48566582	RGR	N	03	035	043	4	3B		0		0						WE	3B	
10	SU48106570	LEY	N	02			1	1	58	-42	58	-33	3B					DR	3B	IMPGH 45
11	SU48206570	LEY	N	05			1	1	44	-56	44	-47	4					DR	4	IMPGH 50
110	SU48506570	LEY	N	05			1	1	50	-50	47	-44	4					DR	4	DR TO 120
14	SU48506570	RGR	N	04	032	042	4	3B		0		0						WE	3B	SPL 42
15	SU48206560	LEY					1	1	54	-46	54	-37	3B					DR	3B	IMPGH 40
16	SU48306560	RGR	NE	05			1	1	44	-56	44	-47	4					DR	4	IMPGH 35
17	SU48406560	PGR	NÉ	05			1	1	29	-71	29	-62	4					DR	4	IMPGH 20
170	SU48406560	PGR	NE	05			1	1	40	-60	37	-54	4					DR	4	DR TO 120
18	SU48506560	PGR	N	05			1	1	31	-69	31	-60	4					DR	4	IMPGH 22
180	SU48506560	PGR	N	05			1	1	44	-56	41	-50	4					ÐR	4	DR TO 120
19	SU48306550	PGR					1	1	58	-42	58	-33	3B					DR	38	IMPGH 40
20	SU48506550	PGR					1	1	34	-66	34	-57	4					DR	4	IMPGH 28
200	SU48506550	PGR	N				1	1	46	-54	43	-48	4					DR	4	DR TO 120

program: ALC011

COMPLETE LIST OF PROFILES 23/03/94 NEWBURY LP SITE 48

page 1

1					MOTTLES	S -	PED			\$1	ONES-		STRUCT	/ SL	BS				
SAMPLE	DEPTH	TEXTURE	COLOUR	COL	ABUN	CONT	COL.	GLE	Y >2	>6	LITH	TOT	CONSIS	rs T	R PO	R IMP	SPL	CALC	
2P	0-15	mcl	10YR32 42						0	0	HR	2							
	15-28	hc1	10YR42 00						0	0	HR	20	WKCSAB	FR N	1				
	28-43	c	10YR52 00	10YR5	6 00 M	2	25Y 52	00 Y	0	٥		Đ	MDCSAB	FR N	1 Y				
i	43-60	С	10YR53 00	75YR5	B 00 M	2	25Y 52	00 Y	0	0		0	MDCAB	FR N	l Y		Υ		
3P	0-24	mcl	10YR32 42						17	0	HR	41							
-	24-50	csl	10YR42 00						16	0	HR	57		١	i				
	50-120	gh	75YR43 00						0	0		0		F	•				
3 Q	0-24	mcl	10YR32 42						17	0	HR	41							
1	24-50	csl	10YR42 00						16	0	HR	57		1	1				
5	50-59	gh (75YR43 00						0	0		0		,)				
4	0-22	mcl	10YR42 00							0	HR	10							
	22-38	c	10YR53 00	10YR5	6 00 C	(OOMMOO	00 Y	0	0	HR	25		F	•		Υ		
•	38-50	С	10YR52 00	10YR5	6 00 M	(OOMMOO	00 Y	0	0	HR	40		F)		Y		IMP STONES 50
4P	0-15		10YR32 42						11	0	HR	29							
ļ	15–58		10YR42 00								HR	55		t	1				
	58-76	mc1	10YR53 00			•	10YR52					10	MDCSAB	FR I	1				
1		С	10YR52 00					Y	0			5		١	1		Υ		STRUCT AS 2P 43-60
	95-110	C	75YR58 Q0	10YR5	3 00 F			Y	0	0	HR	30		f	•		Υ		IMP STONES 110
5	0-28	mcl	10YR41 00	10YR4	6 00 F						HR	10							
	28-37	mcl	10YR53 00						O	0	HR	40		!	1				IMP GRAVEL 37
6	0-28	mcl	10YR42 00								HR	20							
	28-30	mcl	10YR42 52						0	0	HR	40		1	1				IMP GRAVEL 30
6Q	0-28	mcl	10YR42 00						0	0	HR	10							
1	28-30	mcl	10YR42 52						0	0	HR	40		1)				
j	30–120	gh	00ZZ00 00						0	0		0		1)				
9	0-20	mcl	10YR32 42						0	0	HR	3							
	20-35	hc1	10YR42 00						0		HR	5		ľ	1				
,	35-43	С	25Y 51 00					Υ	_		HR	15			•		Υ		
1	43-90	C	25Y 61 00	10YR6	8 00 M			Y	0	0		0		ı	5		Y		
10	0-35	mcl	10YR32 00						0	0	HR	20							
	35-45	mc1	10YR42 00						0	0	HR	60		1	1				IMP GRAVEL 45
11	0-26	mcl	10YR32 00						0	0	HR	20							
•	26-50	gh	10YR54 00						0	0		0		I)				IMP GRAVEL 50
110	0-26	mcl	10YR32 00						0	0	HR	20							
J	26-50	gh	10YR54 00						0	0		0		í)				
	50-120	gh	00ZZ00 00						0	0		0		1)				
7																			

SAMPLE	DEPTH	TEXTURE	COLOUR		MOTTLES ABUN								STRUCT/ CONSIST		x∩p :	TMD S	DI CALC			
	<i>D</i> 2,,	, LATORE	0020011	000	A00/4	CONT	COL.	GEL,		-0	L1 111	101	00/(313)	JIK F	OK .	Ira O	re onec			
14	0-20	mcl	10YR32 0	0					0	0	HŘ	5								
	20-32	mcl	10YR32 5	3					0	0	HR	30		М						
	32-42	mc1	10YR42 0	0 10YR5	6 00 C			Υ	0	0	HR	30		М						
•	42-120	С	25Y 62 0	0 10YR6	M 00 8			Υ	0	0		0		P			Y			
15	0-35	mcl	10YR32 4	2					0	0	HR	20								
	35–40	msl	10YR42 0	0					0	0	HR	60		М				IMP	GRAVEL	40
16	0-20	mcl	10YR31 0	0					0	0	нR	15								
	20-35	mcl	10YR42 0	0					0	0	HR	50		М				IMP	GRAVE	. 35
17	0-15	mcl	10YR31 0	0					0	0	HR	15								
	15-20	m¢1	10YR41 0	0					0	0	HR	50		M				IMP	GRAVEI	_ 20
17Q	0-15	mcl	10YR31 0	0					0	0	HR	15								
	15-20	mcl	10YR41 0	0					0	0	HR	50		М						
	20-120	gh	00ZZ00 0	0					0	0		0		P						
18	0-17	mcl	10YR31 0	0					0	0	HR	15								
	17-22	mcl	75YR31 0	0					0	0	HR	40		М				IMP	GRAVEI	. 22
18Q	0-17	mcl	10YR31 0	0					0	0	HR	15								
	17-22	mcl	75YR31 0	10					0	0	HR	40		М						
	22-120	gh	00ZZ00 0	0					0	0		0		Р						
19	0-20	mcl	10YR32 0						0		HR	10								
	20-35	mcl	10YR42 0						0	0	HR	15		М						
	35–40	നാറി	10YR53 0	00					0	0	HR	40		М				IMP	GRAVE	L 40
20	0-15	mc1	10YR31 0								HR	15								
	15-28	mc1	10YR31 4	1					0	0	HR	50		М				IMP	GRAVE	L 28
20Q	0-15	m¢l	10YR31 C	00					0	0	HR	15								
	15-28	mcl	10YR31 4	11					0	0	HR	50		М						
	28-120	gh	00ZZ00 C	00					0	0		0		Р						