

West Midland Mosses Special Area of Conservation

Evidence Pack

First published August 2022

Natural England Technical Information Note TIN206

West Midland Mosses Special Area of Conservation – Evidence Pack

Anita Wood, Helen Wake and Kathryn McKendrick-Smith



Published August 2022

This report is published by Natural England under the Open Government Licence - OGLv3.0 for public sector information. You are encouraged to use, and reuse, information subject to certain conditions. For details of the licence visit [Copyright](#). Natural England photographs are only available for non-commercial purposes. If any other information such as maps or data cannot be used commercially this will be made clear within the report.

© Natural England 2022

Project details

This report should be cited as: WOOD, A., WAKE, H. and MCKENDRICK-SMITH, K. 2022. *West Midland Mosses Special Area of Conservation – Evidence Pack*. Natural England Technical Information Note. TIN206 Natural England.

Natural England Project manager

Simon Thompson

Author

Anita Wood, Helen Wake and Kathryn McKendrick-Smith

Keywords

Natural England, Nutrient Neutrality, Strategic Solutions, West Midland Mosses SAC

Further information

This report can be downloaded from the Natural England Access to Evidence Catalogue: <http://publications.naturalengland.org.uk/>. For information on Natural England publications contact the Natural England Enquiry Service on 0300 060 3900 or e-mail enquiries@naturalengland.org.uk.

Contents

West Midland Mosses Special Area of Conservation – Evidence Pack	2
Project details	3
Natural England Project manager.....	3
Author	3
Keywords.....	3
1. Site Details	5
2. Reasons for European Designation.....	5
3. Nutrient Pressure and Water Quality	6
4. Additional Information.....	13
List of tables.....	15
List of figures	15
Appendices	16
Bibliography	20
List of abbreviations	20

1. Site Details

From West Midland Mosses Special Area of Conservation Objectives Supplementary Advice:

Lying in the lowlands of the Shropshire, Cheshire and Staffordshire Plain National Character Area, the West Midlands Mosses Special Area of Conservation (SAC) comprises four individual sites: Clarepool Moss, Abbots Moss, Chartley Moss and Wybunbury Moss. These support large basin mires which have developed as quaking bogs, known as 'schwingmoors', together with associated hollows and pools showing various types and stages of mire development, including raised bog on solid peat at two sites. This complexity of habitats gives rise to a diverse assemblage of associated plants and invertebrates of international significance.

The underlying bedrock geology of the area is almost entirely formed of red and brown Triassic sandstones, silts and muds. Glacial activity has affected the whole plain by rounding off hard outcrops of sandstone, creating meltwater channels and lake beds and depositing a variety of materials from boulder clay to marls, sands and gravels. These deposits have in places caused the formation of a number of shallow meres and peat-filled mosses which are present today.

The component sites have been all modified by human activity to a greater or lesser extent, including drainage, peat cutting and nutrient enrichment, generally to the detriment of the natural features.

The restoration of the sites to achieve the best outcomes for the SAC features requires re-naturalisation of the hydrological processes that created them, in terms of both water quality/chemistry and the water supply mechanisms, including groundwater and surface water regimes.

2. Reasons for European Designation

The SAC is designated for the following features:

- H3160 Natural dystrophic lakes and ponds
- H7140 Transition mires and quaking bogs

Links to Conservation Advice:

- [Conservation Objectives](#)
- [Conservation Objectives Supplementary Advice](#)

3. Nutrient Pressure and Water Quality

Nutrient pressure(s) for which the following sites are unfavourable:

Wybunbury Moss:

- Nitrogen

Wybunbury Moss and Abbots Moss:

- Phosphorus

In the Conservation Objectives Supporting Advice for West Midlands Mosses SAC it states for phosphorus '**restore stable nutrient levels appropriate for lake type. Water is normally acidic and poor in available nutrients.**' And for nitrogen '**restore nitrogen concentrations to less than 1.5 mg L⁻¹.**'

Water Quality data is reported against the relevant Site of Special Scientific Interest (SSSI) units within the SAC for Abbots Moss and Wybunbury Moss. Both Chartley Moss and Clarepool Moss are currently assessed as favourable for nutrients.

Abbots Moss

Table 1 – Abbots Moss site attribute with water quality targets

Unit name	SSSI Unit	WQ Target		WQ Monitoring Data ¹		Compliance with target – Pass/Fail	
		TP (µg/l)	TN (µg/l)	TP (µg/l)	TN (µg/l)	TP	TN
Forest Camp	1	10	1.5	All higher than 10 – see details below	All below LOD of 0.002	FAIL	PASS

¹Water Quality monitoring data from: APEM water sampling report (October 2017 – March 2018) surface water sites

Figure 1 – Abbots Moss survey site showing sites 1-3 and 10 (outflow) for surface water sampling





Figure 2 – Surface water Nitrate levels at Abbots Moss Site 1-3 and 10 (outflow)



All surface waters failed the 0.01 mg/l total phosphorus target.

Lilly Pool (sample point 1) had an average total P concentration of 0.88 mg/l (Figure 3). However, this result is somewhat skewed by a very high data point (27/11/17) of 5.16 mg/l. It was reported that this sample was taken after a heavy rainfall event and the sample was very turbid. After this event levels declined to around 0.02 mg/l by the end of March 2018.

Both Round Pool (sample point 2) and Gull Pool (sample point 3) had consistently high total P concentrations with an average of 0.09 mg/l at Round Pool and 0.15 mg/l at Gull Pool.

The outflow stream (sample point 10) from the site had very high total P concentrations over 1.5 mg/l up to December 2017 followed by a decline down to 0.02 mg/l by the end of March 2018.

- Both Gull Pool (site 3) and Round Pool (site 2) show signs of eutrophication with loss of Sphagnum moss from the pool margins and significant blooms of algae. The moderately elevated levels of TP and the very high TP levels recorded after heavy rain highlight Lilly Pool as being at risk of eutrophication and the eventual loss of the remaining areas of dystrophic water at Abbots moss.
- Phosphorus is the key nutrient with respect to the degradation in the habitat quality within the open waters at Abbots moss.

Any nutrients entering the catchment upstream of the locations which are exceeding their nutrient targets have the potential to further add to the current exceedance. Therefore, the whole catchment of Abbots Moss is included within the catchment map.

Wynbunbury Moss

Table 2 – Wynbunbury Moss site water quality targets

Water quality pollutant	SSSI objective/target	Compliance (using WQ monitoring data and/or modelling)	Evidence used to support assessment
Phosphate	The Common Standards Monitoring (2015) standing water habitat phosphorus target for deep dystrophic peak lakes such as the open water at Wynbunbury is 10µg/l (0.01mg/l)	Fails to meet target Total phosphorous concentrations ranged from 0.0209 to 6.51 mg/l. Levels above 1 mg/l were recorded at 5 sites (out of 20 sampled) at the Moss.	Bill Bellamy Associates, 2015. Analysis of Surface Water Quality Samples from Wynbunbury Moss NNR. APEM, 2016. Water Quality Monitoring of

	total phosphorus (soluble + particulate phosphorus)	<p>Concentrations of orthophosphate were above the detection limit of 0.01 mg/l or 10 µg/l in 11 of the 20 samples taken.</p> <p>In a separate six month study, APEM found Total Phosphate concentrations were highly variable ranging from 0.01 mg/l to 1.68 mg/l. See tables below.</p>	Cheshire Meres and Mosses
Nitrogen	The Water Framework Directive (WFD) Groundwater Dependent Terrestrial Ecosystem (GWDTE) threshold value for Wybunbury.Moss is 2mg/l N.	<p>Fails to meet target Observations of nitrate concentration (nitrate-N, mg/l) in the sands and gravels to the north of Wybunbury Moss vary on average between 0.10 mg-N/l and 41.44 mg-N/l</p> <p>In a six month study, APEM found surface water Nitrogen concentrations at Wybunbury Moss ranged from 0.21 mg/l to 14.20 mg/l. An increase in concentrations at most sites occurred during the wetter period between November and February.</p> <p>See tables below.</p>	<p>Environment Agency, 2017. Nitrate Modelling of Wybunbury Moss SSSI wetland catchment SC160010/R.</p> <p>APEM, 2016. Water Quality monitoring of Cheshire Meres and Mosses</p>

The 2012 Natural England CSM assessment concluded unfavourable condition due to presence of negative indicators indicative of enrichment or drying out (*Typha latifolia*); cover of trees / scrub; cover of *Molinia caerulea*; Sphagnum species distribution.

The site was previously very species-rich with records for a large number of rare species notably Rannoch rush (*Scheuchzeria palustris*) and the mosses *Dicranum bergeri* and species characteristic of oligotrophic base-rich groundwater fed fen vegetation. All of these species now believed to be extinct, with major losses of those species characteristic of very low nutrient conditions (EA 2010).

Loss of bryophytes characteristic of low nutrient conditions, particularly from the base-rich areas, and spread of mosses of higher nutrient conditions. Presence of species indicative of nutrient enrichment or pollution as shown in Wheeler et al. (2016).

Any nutrients entering the groundwater or surface water catchment upstream of the locations which are exceeding their nutrient targets have the potential to further add to the current exceedance. Therefore, the whole catchment of Wybunbury Moss is included within the catchment map.

Figure 3 – Total surface water nitrogen and phosphorus (mg/l) concentrations at Wybunbury Moss between August 2015 and January 2016 (APEM 2016)




Table 3 – Wynbunbury Moss total nitrogen (mg/l) concentrations

Sample point	First sample	Last sample	Min (mg-N/l)	Max (mg-N/l)	Average (mg-N/l)
1	27/08/2015	14/01/2016	4.77	14.20	7.04
2	27/08/2015	14/01/2016	1.31	3.09	2.16
3	27/08/2015	14/01/2016	1.09	4.83	2.74
4	27/08/2015	14/01/2016	0.84	1.14	0.96
5	27/08/2015	14/01/2016	1.25	1.93	1.48
6	27/08/2015	14/01/2016	0.43	2.11	1.26
7	27/08/2015	14/01/2016	0.21	3.92	2.62
8	27/08/2015	14/01/2016	1.31	1.59	1.50
9	27/08/2015	14/01/2016	1.64	2.14	1.85

Table 4 – Wynbunbury Moss total phosphorus (mg/l) concentrations

Sample point	First sample	Last sample	Min (mg-P/l)	Max (mg-P/l)	Average (mg-P/l)
1	27/08/2015	14/01/2016	0.052	1.680	0.467
2	27/08/2015	14/01/2016	0.112	0.844	0.423
3	27/08/2015	14/01/2016	0.060	0.439	0.221
4	27/08/2015	14/01/2016	0.01	0.028	0.016
5	27/08/2015	14/01/2016	0.016	0.041	0.025
6	27/08/2015	14/01/2016	0.01	0.106	0.049
7	27/08/2015	14/01/2016	0.01	0.961	0.409


8	27/08/2015	14/01/2016	0.093	0.146	0.115
9	27/08/2015	14/01/2016	0.166	0.429	0.316

Table 5 – Observations of nitrate concentration (nitrate-N, mg/l) in the sands and gravels to the north of Wynbunbury Moss (from Wheeler et al, 2016)

Sample point	First sample	Last sample	Min (mg-N/l)	Max (mg-N/l)	Average (mg-N/l)	n
Wynbunbury Moss – SGA2	03/06/2009	20/01/2010	23.70	28.30	24.86	5
Wynbunbury Moss – SGA3	03/06/2009	10/03/2016	15.50	21.40	18.66	14
Wynbunbury Moss – SGB2	03/06/2009	20/01/2010	37.10	46.00	41.44	5
Wynbunbury Moss – SGB3	03/06/2009	20/01/2010	30.10	36.70	32.44	5
Wynbunbury Moss – SGC2	04/06/2009	20/01/2010	35.20	39.30	37.96	5
Wynbunbury Moss – SGC3	04/06/2009	20/01/2010	0.22	3.88	1.14	5
Wynbunbury Moss – SGD2	04/06/2009	20/01/2010	3.09	9.75	6.71	5
Wynbunbury Moss C 5J65/24	07/07/2003	15/11/2007	13.50	29.30	20.90	4

Wynbunbury Moss Borehole D SJ65/25	10/03/2008	16/08/2016	10.90	39.30	21.10	22
Wynbunbury Moss Lag Fen	06/11/2015	10/03/2016	0.10	0.10	0.10	3

Figure 4 – Location of water quality monitoring sites (from Wheeler et al, 2016)



4. Additional Information

Habitat type impacted by nutrients – Standing Waters

The West Midlands Mosses SAC is legally underpinned by multiple SSSIs. The SSSIs being considered here are Abbots Moss SSSI and Wynbunbury Moss SSSI.

Abbots Moss

SSSI Interest Features include:

- Basin fen (lowland)
- Dystrophic lakes
- Oligotrophic lakes
- Outstanding dragonfly assemblage
- Standing waters

Wynbunbury Moss

SSSI Interest Features:

- Basin fen (lowland) FB
- Quaternary of the Pennines and adjacent areas
- Invert. assemblage W3 permanent wet mire
- IS - Quaternary of the Pennines and adjacent areas

List of tables

Table 1 – Abbots Moss site attribute with water quality targets

Table 2 – Wynbunbury Moss site water quality targets

Table 3 – Wynbunbury Moss total nitrogen (mg/l) concentrations

Table 4 – Wynbunbury Moss total phosphorus (mg/l) concentrations

Table 5 – Observations of nitrate concentration (nitrate-N, mg/l) in the sands and gravels to the north of Wynbunbury Moss (from Wheeler et al, 2016)

List of figures

Figure 1 – Abbots Moss survey site showing sites 1-3 and 10 (outflow) for surface water sampling

Figure 2 – Surface water Nitrate levels at Abbots Moss Site 1-3 and 10 (outflow)


Figure 3 – Total surface water nitrogen and phosphorus (mg/l) concentrations at Wynbunbury Moss between August 2015 and January 2016 (APEM 2016)

Figure 4 - Location of water quality monitoring sites (from Wheeler et al, 2016)

Appendices

Annex 1 – Component SSSIs of West Midlands Mosses SAC (Abbots Moss)

Map of component SSSIs of West Midlands Mosses SAC (Abbots Moss)



European protected sites requiring nutrient neutrality strategic solutions

Scale: 1:10,000

Component SSSIs of West Midlands Mosses SAC (Abbotts Moss)


- Local Authorities
- SSSI subject to nutrient neutrality strategy
- Nutrient neutrality SSSI catchment
- National Parks

Produced by Defra Spatial Data Science
 © Defra 2021, reproduced with the permission of Natural England, <http://www.naturalengland.org.uk/copyright>.
 © Crown Copyright and database rights 2021. Ordnance Survey licence number 100022021.



Annex 2 – Component SSSIs of West Midland Mosses SAC (Wynbunbury Moss)

Map of component SSSIs of West Midland Mosses SAC (Wynbunbury Moss)



European protected sites requiring nutrient neutrality strategic solutions

Scale: 1:10,000

Component SSSIs of West Midlands Mosses SAC (Wybunbury Moss)

- Local Authorities
- SSSI subject to nutrient neutrality strategy
- Nutrient neutrality SSSI catchment
- National Parks

Produced by Defra Spatial Data Science
 © Defra 2021, reproduced with the permission of Natural England, <http://www.naturalengland.org.uk/copyright>.
 © Crown Copyright and database rights 2021. Ordnance Survey licence number 100022021.



Bibliography

APEM, 2016. Water Quality monitoring of Cheshire Meres and Mosses. For Natural England.

BELLAMY, B, 2015. Analysis of surface water quality samples from Wybunbury Moss NNR. For Natural England

ENVIRONMENT AGENCY, 2010. Refining River Basin Planning through targeted investigations on GWDTE . Environment Agency

ENVIRONMENT AGENCY, 2017. Nitrate Modelling of Wybunbury Moss SSSI wetland catchment SC160010/R. Available from:

https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/1036446/Nitrate_modelling_of_Wybunbury_Moss_SSSI_wetland_catchment_-_report.pdf

WHEELER, B., TRATT, R., EADES, P. AND SHAW, S., 2016. Wybunbury Moss: wetland vegetation survey and eco-hydrological assessment 2015. Part 2: Ecohydrological assessment. Final report to Natural England

List of abbreviations

GWDTE – Groundwater Dependent Terrestrial Ecosystem

SAC – Special Area of Conservation

SSSI – Site of Special Scientific Interest

N – Nitrogen

P – Phosphate

TN – Total Nitrogen

TP – Total Phosphate

WFD – Water Framework Directive

Natural England is here to secure a healthy natural environment for people to enjoy, where wildlife is protected and England's traditional landscapes are safeguarded for future generations.

Natural England publications are available as accessible pdfs from www.gov.uk/natural-england.

Should an alternative format of this publication be required, please contact our enquiries line for more information: 0300 060 3900 or email enquiries@naturalengland.org.uk.

Catalogue code: TIN206

This publication is published by Natural England under the Open Government Licence v3.0 for public sector information. You are encouraged to use, and reuse, information subject to certain conditions. For details of the licence visit www.nationalarchives.gov.uk/doc/open-government-licence/version/3.

Please note: Natural England photographs are only available for non-commercial purposes. For information regarding the use of maps or data visit www.gov.uk/how-to-access-natural-english-lands-maps-and-data.

© Natural England 2022

